

A Condition for Finite Blow-up Time for a Volterra Integral Equation

W. MYDLARCZYK

*Institute of Mathematics, University of Wrocław,
Plac Grunwaldzki 2/4, 50-384 Wrocław, Poland*

Submitted by C. Foias

Received August 15, 1991

The paper is devoted to the study of the equation

$$u(x) = \int_0^x (x-s)^{\alpha-1} g(u(s)) ds \quad (\alpha > 0, x > 0),$$

where g is continuous and nondecreasing. Some estimates of u' and the interval of the existence of a nontrivial solution u are given. © 1994 Academic Press, Inc.

1. INTRODUCTION

In this paper we consider the following integral equation

$$u(x) = \int_0^x (x-s)^{\alpha-1} g(u(s)) ds \quad (x > 0, \alpha > 0), \quad (1.1)$$

where

- (i) g is continuous and nondecreasing with $g(0) = 0$ and $g(x) > 0$ for $x > 0$;
- (ii) g satisfies the following generalized Osgood condition ([1, 2, 4]),

$$\int_0^\delta \left(\frac{u}{g(u)} \right)^{1/\alpha} \frac{du}{u} < \infty \quad (\delta > 0).$$

We are interested only in continuous, nonnegative solutions of (1.1). It is clear that $u \equiv 0$ is a solution of (1.1). In [4] it is shown that under the assumptions i and ii there are other, nontrivial, solutions. Moreover, from classical theorems on the existence of the maximal solution for Volterra integral equations (see [3, Chap. II]) and ideas of [1, 5], we have

Remark 1.1. If i and ii are satisfied, then there exists a maximal solu-

tion u of (1.1). It is a nondecreasing function and a unique solution with the property $u(x) > 0$ for $x > 0$.

From now on we deal with a maximal solution u of (1.1). Since it is nondecreasing, we get

$$u(x) = 1/\alpha \int_0^x (x-s)^\alpha dg(u(s)) \quad (x > 0), \quad (1.2)$$

from which follows

Remark 1.2. u is an absolutely continuous function.

Our aim is to estimate $(u^{-1})'$, where u^{-1} is the inverse to u , from which we obtain estimates of the interval $[0, \bar{K})$ of the existence of u . The main results are stated in the following theorem and its corollary.

THEOREM 1.1. *There exist $\gamma_1, \gamma_2 > 0$ such that*

$$\gamma_1 u'(x) \leq \left\{ \int_0^x [u(x) - u(s)]^{\alpha-1} dg(u(s)) \right\}^{1/\alpha} \leq \gamma_2 u'(x),$$

for $0 \leq x \leq \bar{K}$.

If

$$\psi(x) = \left\{ \int_0^x [u(x) - u(s)]^{\alpha-1} dg(u(s)) \right\}^{1/\alpha} \quad (0 < x \leq \bar{K})$$

and $\phi(y) = \psi(u^{-1}(y))$, then from Theorem 1.1 we obtain immediately

COROLLARY 1.1. (i) $\gamma_1 \phi(y) \leq (u^{-1})'(y) \leq \gamma_2 \phi(y)$;

(ii) $\gamma_1 \int_0^\infty \phi(y) dy \leq \bar{K} \leq \gamma_2 \int_0^\infty \phi(y) dy$.

Hence, \bar{K} is finite if and only if $\int_0^\infty \phi(y) dy < \infty$, where $\phi(y) = \left\{ \int_0^y (y-s)^{\alpha-1} dg(s) \right\}^{-1/\alpha}$. The last statement leads to the following theorem.

THEOREM 1.2. *The interval $[0, \bar{K})$ of the existence of u is finite if and only if*

$$\int_0^\infty \left[\frac{s}{g(s)} \right]^{1/\alpha} \frac{ds}{s} < \infty.$$

2. PROOF OF THE THEOREMS

To prove Theorem 1.1 we need two lemmas, in which we consider the cases $0 < \alpha \leq 1$ and $\alpha > 1$, respectively.

Let

$$\Phi(s) = \int_0^s (x-t)^{\alpha-1} dg(u(t)) \quad \text{for } 0 \leq s \leq x.$$

LEMMA 2.1. *If $0 < \alpha \leq 1$, then for $0 \leq x < \bar{K}$*

$$(x-y) \Phi(y) \leq u(x) - u(y) \leq (1/\alpha)(x-y) u'(x). \quad (2.1)$$

Proof. By the Mean Value Theorem we have

$$\alpha(x-s)^{\alpha-1}(x-y) \leq (x-s)^{\alpha} - (y-s)^{\alpha} \quad \text{for } 0 < s < y \leq x. \quad (2.2)$$

Since, by $0 < \alpha \leq 1$, $(y-s)^{\alpha} \geq (x-s)^{\alpha-1}(y-s)$ for $0 < s < y \leq x$, we get

$$(x-s)^{\alpha} - (y-s)^{\alpha} \leq (x-s)^{\alpha-1}(x-y) \quad \text{for } 0 < s < y \leq x. \quad (2.3)$$

Note also that by (1.2)

$$\alpha(u(x) - u(y)) = \int_0^y \{(x-s)^{\alpha} - (y-s)^{\alpha}\} dg(u(s)) + \int_y^x (x-s)^{\alpha} dg(u(s)).$$

Hence using (2.2) and (2.3) we can obtain the required assertion.

COROLLARY 2.1. *If $0 < \alpha \leq 1$, then for $0 \leq x < \bar{K}$*

$$\alpha^{1-\alpha} [u'(x)]^{\alpha} \leq [\psi(x)]^{-\alpha} \leq \alpha^{-1} [u'(x)]^{\alpha}. \quad (2.4)$$

Proof. Since $\alpha - 1 \leq 0$, from the right-hand side inequality of (2.1) it follows that

$$\begin{aligned} [\psi(x)]^{-\alpha} &= \int_0^x [u(x) - u(y)]^{\alpha-1} dg(u(y)) \\ &\geq \alpha^{-(\alpha-1)} [u'(x)]^{\alpha-1} \int_0^x (x-y)^{\alpha-1} dg(u(y)) = \alpha^{1-\alpha} [u'(x)]^{\alpha}. \end{aligned}$$

Noting that $d\Phi(y) = (x-y)^{\alpha-1} dg(u(y))$ and that $\Phi(x) = u'(x)$, and using the left-hand side inequality of (2.1), we obtain

$$\begin{aligned} [\psi(x)]^{-\alpha} &\leq \int_0^x (x-y)^{\alpha-1} \Phi(y)^{\alpha-1} dg(u(y)) \\ &= \int_0^x \Phi(y)^{\alpha-1} d\Phi(y) = (1/\alpha) [u'(x)]^{\alpha}, \end{aligned}$$

which gives the right-hand side inequality of (2.4).

The second lemma concerns the case $\alpha > 1$. The right-hand side inequality is an easy consequence of the Mean Value Theorem. The left-hand side inequality is difficult and depends on an inductive argument.

LEMMA 2.2. *If $\alpha > 1$, then for $0 \leq x < \bar{K}$,*

$$\alpha^{-\alpha} [u'(x)]^\alpha \leq [\psi(x)]^{-\alpha} \leq [u'(x)]^\alpha. \quad (2.5)$$

Proof. We restrict ourself to the left-hand side inequality only.

Let $1 < \alpha = n + \beta$, where $0 < \beta \leq 1$ and n is an integer. Obtaining $u^{(k+1)}$ ($0 \leq k \leq n-1$) from (1.2) and then using the Schwarz inequality, we get

$$[u^{(k+1)}(x)]^2 \leq \frac{\alpha - k}{\alpha - k - 1} u^{(k)}(x) u^{(k+2)}(x) \quad (2.6)$$

for $0 \leq k \leq n-1$ and $0 \leq x < \bar{K}$.

The inequality

$$u''(x) u^{(k)}(x) \leq \frac{\alpha - 1}{\alpha - k} u'(x) u^{(k+1)}(x), \quad 1 \leq k \leq n, \quad (2.7)$$

for $0 \leq x < \bar{K}$ follows from the following inductive arguments. For $k=1$ (2.7) is obvious. Assume (2.7) for $1 \leq k \leq n-1$, then for $k+1$ we have

$$\begin{aligned} u''(x) u^{(k+1)}(x) &= u''(x) u^{(k)}(x) u^{(k+1)}(x) [u^{(k)}(x)]^{-1} \\ &\leq \frac{\alpha - 1}{\alpha - k} u'(x) [u^{(k+1)}(x)]^2 [u^{(k)}(x)]^{-1}. \end{aligned}$$

Now, it suffices to apply (2.6) to get the required assertion.

Since for $0 \leq k \leq n-1$,

$$\begin{aligned} \{[u'(x)]^{k+\beta} u^{(n-k)}(x)\}' &= (k+\beta)[u'(x)]^{k+\beta-1} u''(x) u^{(n-k)}(x) \\ &\quad + [u'(x)]^{k+\beta} u^{(n+1-k)}(x), \end{aligned}$$

from (2.7) we obtain

$$\{[u'(x)]^{k+\beta} u^{(n-k)}(x)\}' \leq \alpha [u'(x)]^{k+\beta} u^{(n+1-k)}(x) \quad (0 \leq k \leq n-1). \quad (2.8)$$

Now we are ready to consider the left-hand side inequality (2.5). We obtain it taking $k=n$ in the inequality

$$\int_0^x \{u(x) - u(y)\}^{k+\beta-1} dg(u(y)) \geq \alpha^{-k+1} c_{n-k} [u'(x)]^{k+\beta-1} u^{(n+1-k)}(x), \quad (2.9)$$

where $c_0 = 1/\alpha$, $c_{n-k} = \{(n+\beta) \cdots (k+\beta)\}^{-1}$ for $0 \leq k \leq n-1$ and $0 \leq x < \bar{K}$, which depends on the following inductive argument.

Let $k=0$. Since $\beta-1 \leq 0$ and u' is nondecreasing, an application of the Mean Value Theorem gives

$$\int_0^x \{u(x) - u(y)\}^{\beta-1} dg(u(y)) \geq \alpha c_n [u'(x)]^{\beta-1} u^{(n+1)}(x).$$

Assume (2.9) for $0 \leq k \leq n-1$. Then for $k+1$ we obtain

$$\begin{aligned} & \int_0^x \{u(x) - u(s)\}^{k+\beta} dg(u(s)) \\ &= (k+\beta) \int_0^x u'(s) \int_0^s \{u(s) - u(t)\}^{k+\beta-1} dg(u(t)) ds \\ &\geq \alpha^{-k+1} (k+\beta) c_{n-k} \int_0^x [u'(s)]^{k+\beta} u^{(n+1-k)}(s) ds. \end{aligned}$$

Since $c_{n-(k+1)} = (k+\beta) c_{n-k}$ and $u^{(n-k)}(0) = 0$ for $0 \leq k \leq n-1$, it suffices to apply the inequality (2.8) to obtain the required assertion.

Proof of Theorem 1.2. First we consider the case $0 < \alpha \leq 1$. An application of the Jensen inequality gives

$$\begin{aligned} & 1/a \int_a^{2a} \left\{ \int_0^s (s-t)^{\alpha-1} dg(t) \right\}^{-1/\alpha} ds \\ &\geq a^{1/\alpha} \left\{ \int_a^{2a} \int_0^s (s-t)^{\alpha-1} dg(t) ds \right\}^{-1/\alpha} \quad (a > 0). \end{aligned} \quad (2.10)$$

Since

$$\int_0^{2a} \int_0^s (s-t)^{\alpha-1} dg(t) ds \leq 1/\alpha \int_0^{2a} (2a-s)^\alpha dg(s) \leq (1/\alpha)(2a)^\alpha g(2a),$$

from (2.10) we get

$$\begin{aligned} & \int_a^{2a} \left\{ \int_0^s (s-t)^{\alpha-1} dg(t) \right\}^{-1/\alpha} ds \geq 2^{-1} (\alpha a)^{1/\alpha} [g(2a)]^{-1/\alpha} \\ &\geq c \int_{2a}^{4a} \left[\frac{s}{g(s)} \right]^{1/\alpha} \frac{ds}{s}, \end{aligned}$$

where $c = 2^{-1} (4^{1/\alpha} - 2^{1/\alpha})^{-1} \alpha^{1/\alpha-1}$. Now it is a simple matter to get

$$\int_0^\infty \phi(y) dy \geq c \int_0^\infty \left[\frac{s}{g(s)} \right]^{1/\alpha} \frac{ds}{s}.$$

The estimate

$$\int_0^s (s-t)^{\alpha-1} dg(t) \geq s^{\alpha-1} g(s)$$

leads to the inequality

$$\int_0^\infty \phi(y) dy \leq \int_0^\infty \left[\frac{s}{g(s)} \right]^{1/\alpha} \frac{ds}{s},$$

which ends the proof of the theorem in the case of $0 < \alpha \leq 1$.

In the case $\alpha > 1$ the required assertion is a simple consequence of the inequality

$$(s/2)^{\alpha-1} g(s/2) \leq \int_0^s (s-t)^{\alpha-1} dg(t) \leq s^{\alpha-1} g(s).$$

ACKNOWLEDGMENT

The author expresses his thanks to the referee for several comments and suggestions helpful in the preparation of the final version of this paper.

REFERENCES

1. P. J. BUSHELL AND W. OKRASINSKI, Uniqueness of solutions for a class of nonlinear Volterra integral equations with convolution kernel, *Math. Proc. Cambridge Philos. Soc.* **106** (1989), 547–552.
2. G. GRIPENBERG, Unique solutions of some Volterra integral equations, *Math. Scand.* **48** (1981), 59–67.
3. R. K. MILLER, "Non-linear Volterra Equations," Benjamin, Menlo Park, 1971.
4. W. MYDLARCZYK, The existence of nontrivial solutions of Volterra equations, *Math. Scand.* **68** (1991), 83–88.
5. W. OKRASINSKI, On a nonlinear Volterra equation, *Math. Methods Appl. Sci.* **18** (1986), 345–350.